



DNS szekvenálás




1. Sanger első módszere (+/- módszer?) (rNTP beépítés és kémiai/enzimes degradáció) 1974:
- Enzimes degradáció, papírkromatográfia majd elektroforézis merőlegesen jellegzetes mintázat, amiből a szekvencia körülményesen, de kikövetkeztethető (ujjlenyomat).




DNS szekvenálás

2. Maxam-Gilbert módszer (részleges kémiai degradáció) 1977.

- Különbségtétel a két-két bázis között:
 - Metilezés dimetil szulfáttal - G 5x gyorsabban metileződik, mint A); Instabil glikozidkötés miatt a nukleobázis melegítésre lehasad semleges pH-n (0.5M HCl, 0°C szinte csak A depurinálódik.)
 - Reakció hidrazinnal (C, T is reagál, de 2M NaCl-ban csak a C!); Nukleobázis elimináció + hidrazon képződés a cukorból.
 - lúgos hidrolízis (0.1M NaOH v. 0.5M piperidin elhasítja a láncot a hiányzó nukleobázisnál.





Frederick Sanger


3. Sanger módszer (terminációs módszer) 1977. Eredeti változat, radioaktív jelölés






A

Radioaktívan jelölt dATP


V = ddATP




Különböző méretű fragmenseket produkál




DNS szekvenálás



DNS szekvenálás




DNS szekvenálás




DNS szekvenálás

ddTTP = **T**



DNS szekvenálás

ddGTP = **G**




Hogyan határozzuk meg a DNS szekvenciát ?

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18

A	1				5				9	10		12					17	
T		2				6					11						16	
G				4				8					13		15			
C			3				7							14				18

Hogyan határozzuk meg a DNS szekvenciát ?


A DNS fragmensek egy nukleotidban különböznek, s elválaszthatóak gél elektroforézissel:




De, e csíkok alapján nem azonosíthatóak a terminális nukleotidok.

DNS szekvenálás


PRINCIPLES OF DNA SEQUENCING BY THE DIDEOXY CHAIN TERMINATION METHOD OF SANGER *ET AL.*




Szekvenálás radioaktívan-jelölt ddNTP-kkel



DNS szekvenálás




DNS szekvenálás



Ugyanaz a módszer, de a primer végén fluoreszcens festék a P^{32} helyett.
Elvileg egy festéssel nem sok különbség. De négy különböző festéssel már igen!


DNS szekvenálás



Ugyanaz a módszer, de a primer végén fluoreszcens festék a P^{32} helyett.
Elvileg egy festéssel nem sok különbség. De négy különböző festéssel már igen!



Modern DNS szekvenálás

Sanger módszer (terminációs módszer) 1987. Prober és mtsai
harmadik változat, fluoreszcens
didezoxi festékek




Modern DNS szekvenálás

Sanger módszer (terminációs módszer) 1987. Prober és mtsai
harmadik változat, fluoreszcens
didezoxi festékek



AUTOMATED DNA SEQUENCING



AUTOMATED DNA SEQUENCING

Using fluorescent terminators or primers (ABI, Pharmacia ALF express, Li-Cor





Fig. 2. Printout.

DNS szekvenálás DNS chipekkel




Hátrány: hibridizációs különbségek és ismétlődések gondjai!

Piroszekvenálás (pirofoszfátképződés mérése)


Pyrosequencing is a nonfluorescence technique that measures the release of inorganic pyrophosphate, which is proportionally converted into visible light by a series of enzymatic reactions (Ronaghi et al. 1996, 1998). Unlike other sequencing approaches that use 3'-modified dNTPs to terminate DNA synthesis, the pyrosequencing assay manipulates DNA polymerase by single addition of dNTPs in limiting amounts. Upon addition of the complementary dNTP, DNA polymerase extends the primer and pauses when it encounters a noncomplementary base. DNA synthesis is reinitiated following the addition of the next complementary dNTP in the dispensing cycle. The light generated by the enzymatic cascade is recorded as a series of peaks called a pyrogram, which corresponds to the order of complementary dNTPs incorporated and reveals the underlying DNA sequence.

The 454 Corporation has recently introduced a whole genome sequencing strategy by integrating pyrosequencing with their PicoTiterPlate (PTP) platform, which has been shown to amplify and image approximately 300,000 PCR templates captured on Sepharose beads (Leamon et al. 2003).

Piroszekvenálás (pirofoszfátképződés mérése)




Piroszekvenálás (pirofoszfátképződés mérése)



Schematic representation of the progress of the enzyme reaction in solid-phase pyrosequencing. The four different nucleotides are added stepwise to the immobilized primed DNA template and the incorporation event is followed using the enzyme ATP sulfurylase and luciferase. After each nucleotide addition, a washing step is performed to allow iterative addition.

Piroszekvenálás (pirofoszfátképződés mérése)



Pyrogram of the raw data obtained from liquid-phase pyrosequencing. Proportional signals are obtained for one, two, three, and four base incorporations. Nucleotide addition, according to the order of nucleotides, is indicated below the pyrogram and the obtained sequence is indicated above the pyrogram.



Workflow

The GS Junior System offers an end-to-end sequencing solution from sample preparation and sequence generation through data analysis. Robust protocols with minimal handling steps make the workflow ideally suited for individual labs. Produce libraries in less than half a day with easy-to-follow sample preparation protocols. Use only general laboratory equipment without the need to purchase tons of additional supplies. Perform overnight sequencing and data processing with a quick 10-hour instrument run time. Go from sequence data to publishable result with straightforward tools for de novo assembly, mapping and amplicon variant analysis.

How It Works?

1. Sample Input and Fragmentation

The GS Junior System supports the sequencing of samples from a wide variety of starting materials including genomic DNA, PCR products, BACs, and cDNA. Samples such as genomic DNA and BACs are randomly fragmented into small, 300- to 800-basepair pieces. For smaller samples, such as small non-coding RNA or PCR amplicons, fragmentation is not required. Instead, PCR products amplified using Genome Sequencer fusion primers may be immobilized onto DNA capture beads and clonally amplified as shown below under "One Fragment = One Bead".

2. Library Preparation

Using a series of standard molecular biology techniques, short DNA adaptors are added to each library fragment. These adaptors are then used in subsequent quantification, amplification, and sequencing steps.

3. One Fragment = One Bead

The single-stranded DNA library is immobilized onto specifically designed DNA Capture Beads. Each bead carries a unique single-stranded DNA library fragment. The bead-bound library is emulsified with amplification reagents in a water-in-oil mixture resulting in microreactors containing just one bead with one unique sample-library fragment.

4. emPCR (Emulsion PCR) Amplification


Each unique sample library fragment is clonally amplified within its own microreactor, excluding competing or contaminating sequences. Amplification of the entire fragment collection is carried out in parallel; for each fragment, this produces several million copies of the original fragment per bead. Subsequently, the emulsions are broken to facilitate collection of the amplified fragments bound to their specific beads.

5. One Bead = One Read

The clonally amplified fragments are enriched and loaded onto a PicoTiterPlate device for sequencing. The diameter of the PicoTiterPlate wells allows for only one bead per well. After addition of sequencing enzymes and reagents, the fluidics subsystem of the Genome Sequencer System serially flows nucleotides in a fixed order (i.e. first T, then A, and so on) across the hundreds of thousands of wells containing one bead each. Addition of one (or more) nucleotide(s) complementary to the template strand results in a chemiluminescent signal recorded by the CCD camera of the Genome Sequencer System. The intensity of the resulting signal is proportional to the number of bases incorporated.

6. Data Analysis

The combination of signal intensity and positional information generated across the PicoTiterPlate device allows the software to determine the sequence of 100,000 individual reads per 10-hour instrument run simultaneously. For sequencing data analysis, three different bioinformatics tools are supplied that readily support the following applications: de novo genome assembly up to 3 Gb; resequencing/mapping genomes of any size; and amplicon variant detection by comparison with a known reference sequence.



TOTAL GENOME SEQUENCING PROJECTS




Fig. 1. A circular representation of the *H. influenzae* Rd chromosome illustrating the location of each

28,984 sequencing reactions

84% success

Ave. read length 485 bp


Genome size 1.8301 Mb

Predicted ORFs 1743

Unassigned 42%

No. of authors 39

Haemophilus influenzae
TIGR (1995)



Box 7.1 *Estimates of the required size of genomic libraries*

Organism	Genome size	Vector type	Insert size	P	Library size
Bacterium	4×10^6 bases	plasmid	4 kb	0.99	4.6×10^3
		lambda replacement	18 kb	0.99	1.0×10^3
		cosmid	40 kb	0.99	458
		BAC	300 kb	0.99	59
Mammal	3×10^9 bases	plasmid	4 kb	0.99	3.5×10^6
		lambda replacement	18 kb	0.99	7.7×10^5
		cosmid	40 kb	0.99	3.5×10^5
		BAC	300 kb	0.99	4.6×10^4

The values shown for the genome sizes of bacteria and mammals are examples for the purpose of this calculation. The actual genome sizes vary quite widely from one organism to another. The insert sizes for specific vectors will also vary.

Table 1–1 Some Genomes That Have Been Completely Sequenced

SPECIES	SPECIAL FEATURES	HABITAT	GENOME SIZE (1000s OF NUCLEOTIDE PAIRS PER HAPLOID GENOME)	ESTIMATED NUMBER OF GENES CODING FOR PROTEINS
BACTERIA				
<i>Mycoplasma genitalium</i>	has one of the smallest of all known cell genomes	human genital tract	580	468
<i>Synechocystis</i> sp.	photosynthetic, oxygen-generating (cyanobacterium)	lakes and streams	3573	3168
<i>Escherichia coli</i>	laboratory favorite	human gut	4639	4289
<i>Helicobacter pylori</i>	causes stomach ulcers and predisposes to stomach cancer	human stomach	1667	1590
<i>Bacillus anthracis</i>	causes anthrax	soil	5227	5634
<i>Aquifex aeolicus</i>	lithotrophic; lives at high temperatures	hydrothermal vents	1551	1544
<i>Streptomyces coelicolor</i>	source of antibiotics; giant genome	soil	8667	7825
<i>Treponema pallidum</i>	spirochete; causes syphilis	human tissues	1138	1041
<i>Rickettsia prowazekii</i>	bacterium most closely related to mitochondria; causes typhus	lice and humans (intracellular parasite)	1111	834
<i>Thermotoga maritima</i>	organotrophic; lives at very high temperatures	hydrothermal vents	1860	1877

Genome size and gene number vary between strains of a single species, especially for bacteria and archaea. The table shows data for particular strains that have been sequenced. For eucaryotes, many genes can give rise to several alternative variant proteins, so that the total number of proteins specified by the genome is substantially greater than the number of genes.

Table 1–1 Some Genomes That Have Been Completely Sequenced

SPECIES	SPECIAL FEATURES	HABITAT	GENOME SIZE (1000s OF NUCLEOTIDE PAIRS PER HAPLOID GENOME)	ESTIMATED NUMBER OF GENES CODING FOR PROTEINS
ARCHAEA				
<i>Methanococcus jannaschii</i>	lithotrophic, anaerobic, methane-producing	hydrothermal vents	1664	1750
<i>Archaeoglobus fulgidus</i>	lithotrophic or organotrophic, anaerobic, sulfate-reducing	hydrothermal vents	2178	2493
<i>Nanoarchaeum equitans</i>	smallest known archaean; anaerobic; parasitic on another, larger archaean	hydrothermal and volcanic hot vents	491	552
EUCARYOTES				
<i>Saccharomyces cerevisiae</i> (budding yeast)	minimal model eucaryote	grape skins, beer	12,069	~6300
<i>Arabidopsis thaliana</i> (Thale cress)	model organism for flowering plants	soil and air	~142,000	~26,000
<i>Caenorhabditis elegans</i> (nematode worm)	simple animal with perfectly predictable development	soil	~97,000	~20,000
<i>Drosophila melanogaster</i> (fruit fly)	key to the genetics of animal development	rotting fruit	~137,000	~14,000
<i>Homo sapiens</i> (human)	most intensively studied mammal	houses	~3,200,000	~24,000

Genome size and gene number vary between strains of a single species, especially for bacteria and archaea. The table shows data for particular strains that have been sequenced. For eucaryotes, many genes can give rise to several alternative variant proteins, so that the total number of proteins specified by the genome is substantially greater than the number of genes.



Phage λ
50 kb
2 pages



Escherichia coli
(bacteria)
4.7 Mb
200 pages



Saccharomyces cerevisiae
(yeast)
12.5 Mb
500 pages



Caenorhabditis elegans
(nematode)
Arabidopsis thaliana
(plant)
100 Mb
3 volumes












Drosophila melanogaster
(fruit fly)
165 Mb
5 volumes











Human being
3000 Mb
80 volumes











25 kb per page
1500 pages
per volume
(2 inches thick)

Amoeba dubia	670 000 000 000	
Amoeba proteus	290 000 000 000	
Lilium longiflorum	90 000 000 000	
Pinus resinosa	68 000 000 000	
Triturus cristatus	20 600 000 000	
Allium cepa	18 000 000 000	
Podisma pedestris	18 000 000 000	
Paramecium caudatum	8 600 000 000	
Bufo bufo	6 900 000 000	

Zea mays	5 000 000 000	
Macropus giganteus	4 154 800 000	
Bos taurus	3 651 500 000	
Pongo pygmaeus	3 607 100 000	
Pan troglodytes	3 577 500 000	
Macaca mulatta	3 543 000 000	
Gorilla gorilla	3 523 200 000	
Papio hamadryas	3 478 800 000	
Mus musculus	3 454 200 000	
Homo sapiens	3 400 000 000	

Caiman crocodylus	2 600 000 000	
Parascaris equorum	2 500 000 000	
Microtus pennsylvanicus	2 477 100 000	
Boa constrictor	2 100 000 000	
Danio rerio	1 900 000 000	
Aplysia californica	1 800 000 000	
Python reticulatus	1 700 000 000	
Cyprinus carpio	1 700 000 000	

Escherichia coli	4 639 221	
Mycobacterium tuberculosis	4 397 000	
Mycobacterium leprae	2 800 000	
Haemophilus influenzae	1 830 137	
Helicobacter pylori	1 667 867	
Methanococcus jannaschii	1 664 974	
Borrelia garinii	953 000	
Mycoplasma genitalium	580 000	

PubMed home - Microsoft Internet Explorer

Fájl Szerkesztés Nézet Kedvencek Eszközök Súgó

Vissza Keresés Kedvencek

Cím <http://www.ncbi.nlm.nih.gov/pubmed/> Ugrás Hivatkozások

NCBI Resources How To My NCBI Sign In

PubMed
U.S. National
National Institute of Health

Search: PubMed Limits Advanced search Help

Search Clear

All Resources
Literature
DNA & RNA
Proteins
Sequence Analysis
Genes & Expression
Genomes & Maps
Domains & Structures
Genetics & Medicine
Taxonomy
Data & Software
Training & Tutorials
Homology
Small Molecules
Variation

Database of Genomic Structural Variation (dbVar)
Genome
Genome Project
Genome Workbench
Influenza Virus
Map Viewer
Nucleotide
PopSet
ProSplign
Sequence Read Archive (SRA)
Splign
Trace Archive
UniSTS
All Genomes & Maps Resources...

Tool for searching, viewing and downloading genomic maps and related data

million citations for
and life science journals.
text articles from PubMed

Using PubMed
PubMed Quick Start
New and Noteworthy
PubMed Tutorials
Full Text Articles
PubMed FAQs

Tools
on Matcher
on Matcher
ies
ic Queries

More Resources
MeSH Database
Journals Database
Clinical Trials
E-Utilities
LinkOut

Map Viewer - Microsoft Internet Explorer

Fájl Szerkesztés Nézet Kedvencek Eszközök Súgó

Vissza

Keresés

Kedvencek

Cím <http://www.ncbi.nlm.nih.gov/mapview/> Ugrás Hivatkozások

NCBI Home GenBank BLAST

Map Viewer Home Help

The Map Viewer provides a wide variety of genome mapping and sequencing data. [More..](#)

Search

Search: Select Group or Organism

for: Select Group or Organism

Tools Le

News

Human

Annotati

human g

An annota

(NCBI Bu

Related

Small Genomes

Select Group or Organism

-- Groups --

Plants

Monocots

Eudicots

Fungi

-- Organisms --

-- Primates ---

Homo sapiens

Macaca mulatta

Pan troglodytes

-- Rodents ---

Mus musculus

Rattus norvegicus

-- Monotremes ---

Ornithorhynchus anatinus

-- Marsupials ---

Monodelphis domestica

-- Other Mammals ---

Bos taurus

Bubalus bubalis

Canis lupus familiaris

Equus caballus

Felis catus

Ovis aries

Sus scrofa

-- Other Vertebrates ---

NCBI Danio rerio

NCBI Gallus gallus

NCBI Taeniopygia guttata

Genome Browser agreement

Genome Biology

Taxonomy

Entrez (Global Query)

BLAST

Map Viewer FTP

Bacteria

Organelles

Viruses

Vertebrates (17)

Mammals (14)

Primates (3)

Scientific name	Common name	Build	Tools
<i>Homo sapiens</i>	human	Build 37.1	<div><div>Q</div><div>B</div><div>R</div><div>C</div><div>G</div></div>

Map Viewer - Microsoft Internet Explorer

Fájl Szerkesztés Nézet Kedvencek Eszközök Súgó

Vissza Vissza Vissza Vissza Vissza Keresés Kedvencek

Cím <http://www.ncbi.nlm.nih.gov/mapview/>

NCBI Home GenBank BLAST

Map Viewer Home

The Map Viewer provides a wide variety of genome mapping and sequencing

Search

Search:

for:

Go

Tools Legend

- Search or Browse the Genome
- BLAST
- Clone Finder
- Go to region on a chromosome
- Genome Resources page

News

Vertebrates

Mammals

Primates



NCBI Map Viewer

PubMed

Nucleotide

Protein

Genome

Gene

Structure

Search for

on chromosome(s)

assembly

All

Find

Map Viewer

[Map Viewer Home](#)

[Map Viewer Help](#)

[Human Maps Help](#)

[Release Notes](#)

NCBI Resources

[Genome Project](#)

[TaxPlot](#)


[Consensus CoDing
Sequence \(CCDS\)](#)

[Human Genome
Resources](#)

[NCBI Handbook](#)

Homo sapiens (human) genome view

[Build 36.2 statistics](#) [Switch to previous build](#)



Human genome
overview page (Build
36.2)
Human genome
overview page (Build
35.1)

[Map Viewer Home](#)

[Map Viewer Help](#)
[Human Maps Help](#)
[FTP](#)

[Data As Table View](#)

[Maps & Options](#)


[Compress Map](#) ☐

Region Shown:

☐ out
☒ zoom
☐ in

You are here:

[Ideogram](#)



☒ default
☐ master

[Homo sapiens \(human\)](#) [Build 36.2 \(Current\)](#)

[BLAST The Human Genome](#)

Chromosome: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [11](#) [12](#) [13](#) [14](#) [15](#) [16](#) [17](#) [18](#) [19](#) [20](#) [21](#) [22](#) [X] Y MT

Master Map: [Genes On Sequence](#)



[Summary of Maps](#)



[Maps & Options](#)

Region Displayed: 0-155M bp

[Download/View Sequence/Evidence](#)

Ideogram	Contig	Hs Uni	Genes_seq	Symbol	Links	E	Cyto	Description
Xp22.33	NT_086925	Hs.350927		VCX-C	sv pr dl ev mm hm	SNP	best RefSeq Xp22	variably charged X-C
Xp22.32	NT_078115	Hs.522584		AP1S2	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xp22.2		adaptor-related protein co
Xp22.31	NT_028413	Hs.495710		MAGEB6	OMIM HGNC sv pr dl ev mm hm	CCDS SNP	best RefSeq Xp21.3	melanoma antigen family I
Xp22.2	NT_086929	Hs.495755		SC4MOP	HGNC sv dl ev mm		best RefSeq Xp21.1	sterol-C4-methyl oxidase
Xp22.13	NT_011757	Hs.28491		UBE1	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xp11.23		ubiquitin-activating enzym
Xp22.12	NT_079573	Hs.441664		CLCN5	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xp11.23-p11.22		chloride channel 5 (nephr
Xp21.3	NT_086939	Hs.533273		GNL3L	HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xp11.22		guanine nucleotide binding
Xp21.2	NT_011638	Hs.522632		APEX2	HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xp11.21		APEX nuclease (apurinic/
Xp21.1	NT_011630	Hs.301404		LOC260337	sv dl ev mm		best RefSeq Xq13.2	zinc finger protein Np97 p
Xq11.2		Hs.533282		BMP2KL	HGNC sv pr dl ev mm hm	SNP	protein Xq13.2	BMP2 inducible kinase-li
Xq11.1		Hs.446628		COX7B	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xq21.1		cytochrome c oxidase sub
Xq12		Hs.529901		LOC644504	sv pr dl ev mm hm	SNP	protein Xq22.3	similar to chondroitin beta
Xq13.1		Hs.17501		PSMD10	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xq22.3		proteasome (prosome, ma
Xq13.2		Hs.522632		LRCH2	HGNC sv pr dl ev mm hm sts	SNP	best RefSeq Xq23	leucine-rich repeats and c
Xq13.3		Hs.496622		LOC727968	sv pr dl ev mm hm		protein Xq23	hypothetical protein LOC
Xq21.1		Hs.632282		PLAC1	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xq26		placenta-specific 1
Xq21.2		Hs.435369		SLITRK4	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xq27.3		SLIT and NTRK-like fam
Xq21.3		Hs.380118		MAGEA12	OMIM HGNC sv pr dl ev mm hm sts CCDS SNP	best RefSeq Xq28		melanoma antigen family
Xq22.1		Hs.460960		LOC728749	sv dl ev mm hm	mRNA Xq28		similar to cancer/testis ant
Xq22.2		Hs.522805		H2AFB2	HGNC sv pr dl ev mm hm	CCDS	best RefSeq Xq28	H2A histone family, meml
Xq22.3		Hs.521						
Xq23		Hs.195464						
Xq24		Hs.534404						
Xq25		Hs.6551						
Xq26		Hs.74576						
Xq27								
Xq28								

Human genome overview page (Build 36.2)
Human genome overview page (Build 35.1)
[Map Viewer Home](#)
[Map Viewer Help](#)
[Human Maps Help](#)
[FTP](#)
[Data As Table View](#)
Maps & Options
[Compress Map](#) ☐
Region Shown:
29,900K
34,900K

You are here:
Ideogram

☒ default
☐ master



MAGEB4	+	OMIM	HGNC	sv pr dl ev mm hm	CCDS	SNP	best RefSeq	Xp21.3	melanoma antigen
MAGEB1	+	OMIM	HGNC	sv pr dl ev mm hm sts	CCDS	SNP	best RefSeq	Xp21.3	melanoma antigen
NR0B1	+	OMIM	HGNC	sv pr dl ev mm hm sts	CCDS	SNP	best RefSeq	Xp21.3-p21.2	nuclear receptor
CXorf21	+		HGNC	sv pr dl ev mm hm	CCDS	SNP	best RefSeq	Xp21.2	chromosome X
LOC652904	+			sv pr dl ev mm hm		SNP	protein	Xp21.2	similar to Cyclin
LOC392437	+			sv dl ev mm	sts		protein	Xp21.2	hypothetical LOC
GK	+	OMIM	HGNC	sv pr dl ev mm hm sts	CCDS	SNP	best RefSeq	Xp21.3	glycerol kinase
MAP3K7IP3	+	OMIM	HGNC	sv pr dl ev mm hm sts	CCDS	SNP	best RefSeq	Xp21.2	mitogen-activated
FTHL17	+	OMIM	HGNC	sv pr dl ev mm hm sts	CCDS	SNP	best RefSeq	Xp21	ferritin, heavy p
DMD	+	OMIM	HGNC	sv pr dl ev mm hm sts	CCDS	SNP	best RefSeq	Xp21.2	dystrophin (mu
TBCAP1	+			sv dl ev mm			best RefSeq	Xp21	tubulin-specific
LOC646506	+			sv dl ev mm			protein	Xp21.1	similar to Lysos
FAM47A	+		HGNC	sv pr dl ev mm hm		SNP	best RefSeq	Xp21.1	family with sequ
LOC139431	+			sv dl ev mm			protein	Xp21.1	similar to ferritin
LOC392439	+			sv dl ev mm			protein	Xp21.1	similar to Splice
TMEM47	+		HGNC	sv pr dl ev mm hm	CCDS	SNP	best RefSeq	Xp21.4	transmembrane


Official Symbol	DMD	provided by HGNC	Genomic context Bibliography Interactions General gene information General protein information Reference Sequences Related Sequences Additional Links
Official Full Name	dystrophin (muscular dystrophy, Duchenne and Becker types)	provided by HGNC	
Primary source	HGNC:2928		
Locus tag	GS1-19024.1		
See related	Ensembl:ENSG00000198947 ; HPRD:02303 ; MIM:300377		
Gene type	protein coding		▼ Links Explain Order cDNA clone Books Conserved Domains Genome GEO Profiles HomoloGene Map Viewer CoreNucleotide Nucleotide OMIM Full text in PMC Probe Protein PubMed PubMed (GeneRIF) SNP SNP: Genotype SNP: GeneView Taxonomy UniSTS AceView CCDS DMD.html Ensembl Evidence Viewer GDB GeneTests for MIM: 300377 GeneTests for MIM: 310200 HGMD HGNC HPRD KEGG Leiden Muscular Dystrophy pages MGC ModelMaker UniGene LinkOut
RefSeq status	Reviewed		
Organism	Homo sapiens		
Lineage	<i>Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo</i>		
Also known as	BMD; CMD3B; DXS142; DXS164; DXS206; DXS230; DXS239; DXS268; DXS269; DXS270; DXS272		
Summary	<p>The dystrophin gene is the largest gene found in nature, measuring 2.4 Mb. The gene was identified through a positional cloning approach, targeted at the isolation of the gene responsible for Duchenne (DMD) and Becker (BMD) Muscular Dystrophies. DMD is a recessive, fatal, X-linked disorder occurring at a frequency of about 1 in 3,500 new-born males. BMD is a milder allelic form. In general, DMD patients carry mutations which cause premature translation termination (nonsense or frame shift mutations), while in BMD patients dystrophin is reduced either in molecular weight (derived from in-frame deletions) or in expression level. The dystrophin gene is highly complex, containing at least eight independent, tissue-specific promoters and two polyA-addition sites. Furthermore, dystrophin RNA is differentially spliced, producing a range of different transcripts, encoding a large set of protein isoforms. Dystrophin (as encoded by the Dp427 transcripts) is a large, rod-like cytoskeletal protein which is found at the inner surface of muscle fibers. Dystrophin is part of the dystrophin-glycoprotein complex (DGC), which bridges the inner cytoskeleton (F-actin) and the extra-cellular matrix.</p>		

Genomic regions, transcripts, and products
↑ ?

(minus strand) Go to [reference sequence details](#)

There is an enormous amount of information here.

NC_000023.9






Figure 1. A: Genomic organisation of the dystrophin gene, located in Xp21. The black vertical lines represent the 79 exons of the dystrophin gene distributed over about 2.5 million bases. The arrows indicate the various promoters: in particular are brain (B), muscle (M), and Purkinje (P) promoters; R, B₃, S, and G represent the Dp260 (retinal), Dp140 (brain), Dp116 (Schwann cells), and Dp71 (general) promoters. B: The domain composition of the various dystrophin proteins is indicated. The amino-terminal domain is followed by the spectrin like domain, the cysteine rich, and the carboxy-terminal domain.

6. Feladat:

A Duchenne-féle izomsorvadásos betegségért felelős DMD gén az X-kromoszóma 21-es lókuszában található. A betegség általában egy-egy kódoló régió hiányának köszönhető, aminek következtében a gén által kódolt disztrofin fehérje nem termelődik, ezért az izomsejtek fokozatosan elhalnak. A humán genom projektnek köszönhetően a gén teljes szekvenciája ismert. Szeretnénk kideríteni, hogy a gén 1. exonja (szekvencia a következő oldalon) megvan-e a páciens DNS-ében? Ehhez szükség lenne az 1. exon szekvenciájából egy olyan 30 bázispárból álló részletre, mely egyedi a humán genomban. Ez annyit jelent, hogy a humán genomban a leghosszabb nemspecifikus kötődés(ek)re jellemző olvadáspont(ok) legalább 15°C-kal legyen(ek) kisebb(ek) mint a 30 bázispáros marker olvadáspontja.

Keressen egy ilyen markert és szekvenciáját, valamint jellemzését küldje el e-mailben a gyurcsik@chem.u-szeged.hu címre! A file neve a monogramja és egy utána írt 6.doc legyen.

A DMD gén 1. exonjának DNS szekvenciája.

**TCCTGGCATCAGTTACTGTGTTGACTCACTCAGTGTTGGGATCACTCAC
TTTCCCCCTACAGGACTCAGATCTGGGAGGCAATTACCTTCGGAGAAAA
ACGAATAGGAAAACTGAAGTGTTACTTTTTTTTAAAGCTGCTGAAGTTT
GTTGGTTTCTCATTGTTTTTTAAGCCTACTGGAGCAATAAAGTTTGAAGA
ACTTTTACCAGGTTTTTTTTTATCGCTGCCTTGATATACACTTTTCAAAA
TGCTTTGGTGGGAAGAAGTAGAGGACTGTT**

PubMed home - Microsoft Internet Explorer

Fájl Szerkesztés Nézet Kedvencek Eszközök Súgó

Vissza

Keresés

Kedvencek

Cím <http://www.ncbi.nlm.nih.gov/pubmed/>

Ugrás

Hivatkozások

NCBI

Resources

How To

My NCBI

Sign In

PubMed

U.S. National Institutes of Health

Search: PubMed

Limits Advanced search Help

Search Clear

All Resources

Literature

DNA & RNA

Proteins

Sequence Analysis

Genes & Expression

Genomes & Maps

Domains & Structures

Genetics & Medicine

Taxonomy

Data & Software

Training & Tutorials

Homology

Small Molecules

Variation

BLAST

BLAST (Stand-alone)

BLAST Link (BLink)

GenBank

Genome Workbench

Influenza Virus

Primer-BLAST

ProSplign

Splign

All Sequence Analysis Resources...

PubMed

More than 19 million citations for MEDLINE and life science journals. Links to full-text articles from PubMed web sites.

Using PubMed

[PubMed Quick Start](#)

[New and Noteworthy](#)

[PubMed Tutorials](#)

[Full Text Articles](#)

[PubMed FAQs](#)

PubMed Tools

[Single Citation Matcher](#)

[Batch Citation Matcher](#)

[Clinical Queries](#)

[Topic-Specific Queries](#)

More Resources

[MeSH Database](#)

[Journals Database](#)

[Clinical Trials](#)

[E-Utilities](#)

[LinkOut](#)

BLAST: Basic Local Alignment Search Tool - Microsoft Internet Explorer

Fájl Szerkesztés Nézet Kedvencek Eszközök Súgó

Vissza

Keresés

Kedvencek

Cím <http://blast.ncbi.nlm.nih.gov/Blast.cgi> Ugrás Hivatkozások

BLAST

Basic Local Alignment Search Tool

HomeRecent ResultsSaved StrategiesHelp

My NCBI
[Sign In] [Register]

NCBI/ BLAST Home

BLAST finds regions of similarity between biological sequences. [more...](#)

New Aligning Multiple Protein Sequences? Try the [COBALT Multiple Alignment Tool](#). Go

BLAST Assembled Genomes

Choose a species genome to search, or [list all genomic BLAST databases](#).

[Human](#)
[Mouse](#)
[Rat](#)
[Arabidopsis thaliana](#)

[Oryza sativa](#)
[Bos taurus](#)
[Danio rerio](#)
[Drosophila melanogaster](#)

[Gallus gallus](#)
[Pan troglodytes](#)
[Microbes](#)
[Apis mellifera](#)

Basic BLAST

Choose a BLAST program to run.

nucleotide blast	Search a nucleotide database using a nucleotide query <i>Algorithms: blastn, megablast, discontinuous megablast</i>
protein blast	Search protein database using a protein query <i>Algorithms: blastp, psi-blast, phi-blast</i>
blastx	Search protein database using a translated nucleotide query
tblastn	Search translated nucleotide database using a protein query
tblastx	Search translated nucleotide database using a translated nucleotide query

News

[COBALT improvements](#)

A COBALT multiple sequence alignment can now be downloaded to a local file.
Thu, 21 Jan 2010 17:00:00 EST

[More BLAST news...](#)

Tip of the Day

[Use Genomic BLAST to see the genomic context](#)

If you are interested in the evolution of a particular gene or gene family it is often interesting to examine the intro-exon structure even across species.

[More tips...](#)

Nucleotide BLAST: Search nucleotide databases using a nucleotide query - Microsoft Internet Explorer

Fájl Szerkesztés Nézet Kedvencek Eszközök Súgó

Vissza Keresés Kedvencek

Cím http://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&BLAST_PROGRAMS=megaBlast&PAGE_TYPE=BlastSearch&SHOW_DEFAULTS=on&LINK_LOC=blasthome Ugrás Hivatkozások

BLAST Basic Local Alignment Search Tool

Home Recent Results Saved Strategies Help

My NCBI [Sign In] [Register]

NCBI/ BLAST/ blastn suite

blastn blastp blastx tblastn tblastx

Enter Query Sequence BLASTN programs search nucleotide databases using a nucleotide query. [more...](#) [Reset page](#) [Bookmark](#)

Enter accession number, gi, or FASTA sequence [Clear](#)

TCCTGGCATCAGTTACTGTGTTGACTCACT

Query subrange

From

To

Or, upload file [Tallózás...](#)

Job Title

Enter a descriptive title for your BLAST search

☐ Align two or more sequences

Choose Search Set

Database ☒ Human genomic + transcript ☐ Mouse genomic + transcript ☐ Others (nr etc.):
Human genomic plus transcript (Human G+T)

Exclude ☐ Models (XM/XP) ☐ Uncultured/environmental sample sequences

Optional Entrez Query

Optional Enter an Entrez query to limit search


Program Selection

Optimize for ☒ Highly similar sequences (megablast) ☐ More dissimilar sequences (discontiguous megablast) ☐ Somewhat similar sequences (blastn)

Choose a BLAST algorithm

Distribution of 1799 Blast Hits on the Query Sequence

Mouse-over to show define and scores, click to show alignments




▼ Descriptions

Legend for links to other resources: [U](#) UniGene [E](#) GEO [G](#) Gene [S](#) Structure [M](#) Map Viewer

Sequences producing significant alignments:

Accession	Description	Max score	Total score	Query coverage	E value	Max ident	Links
Transcripts							
NM_004006.1	Homo sapiens dystrophin (DMD), transcript variant Dp427m, mRNA	60.0	60.0	100%	6e-08	100%	G M
Genomic sequences [show first]							
NT_167197.1	Homo sapiens chromosome X genomic contig, GRCh37 reference pri	60.0	518	100%	6e-08	100%	
NW_001842360.1	Homo sapiens chromosome X genomic contig, alternate assembly (b	60.0	409	100%	6e-08	100%	
NT_023133.13	Homo sapiens chromosome 5 genomic contig, GRCh37 reference pri	36.2	498	90%	0.91	100%	
NW_001838960.2	Homo sapiens chromosome 5 genomic contig, alternate assembly (b	36.2	36.2	60%	0.91	100%	
NT_011651.17	Homo sapiens chromosome X genomic contig, GRCh37 reference pri	34.2	543	96%	3.6	100%	
NT_079573.4	Homo sapiens chromosome X genomic contig, GRCh37 reference pri	34.2	171	93%	3.6	100%	
NT_011387.8	Homo sapiens chromosome 20 genomic contig, GRCh37 reference p	34.2	543	100%	3.6	100%	
NT_030059.13	Homo sapiens chromosome 10 genomic contig, GRCh37 reference p	34.2	1340	96%	3.6	100%	
NT_005403.17	Homo sapiens chromosome 2 genomic contig, GRCh37 reference pri	34.2	1443	100%	3.6	100%	
NW_001842386.2	Homo sapiens chromosome X genomic contig, alternate assembly (b	34.2	220	80%	3.6	100%	
NW_001842361.2	Homo sapiens chromosome X genomic contig, alternate assembly (b	34.2	145	93%	3.6	100%	
NW_001838652.1	Homo sapiens chromosome 20 genomic contig, alternate assembly (34.2	409	100%	3.6	100%	
NW_001837986.1	Homo sapiens chromosome 10 genomic contig, alternate assembly (34.2	276	96%	3.6	100%	
NW_001838860.1	Homo sapiens chromosome 2 genomic contig, alternate assembly (b	34.2	434	100%	3.6	100%	
NT_025028.14	Homo sapiens chromosome 18 genomic contig, GRCh37 reference p	32.2	300	96%	14	100%	
NT_026437.12	Homo sapiens chromosome 14 genomic contig, GRCh37 reference p	32.2	1193	100%	14	100%	
NT_029419.12	Homo sapiens chromosome 12 genomic contig, GRCh37 reference p	32.2	1352	100%	14	100%	
NT_009237.18	Homo sapiens chromosome 11 genomic contig, GRCh37 reference p	32.2	589	96%	14	100%	
NT_008470.19	Homo sapiens chromosome 9 genomic contig, GRCh37 reference pri	32.2	997	96%	14	100%	

> [ref|NT_023133.13|](#)  Homo sapiens chromosome 5 genomic contig, GRCh37 reference primary assembly
Length=25716533

Sort alignments for this subject sequence by:

E value [Score](#) [Percent identity](#)

[Query start position](#) [Subject start position](#)

Features in this part of subject sequence:
[hypothetical protein LOC375484](#)

Score = 36.2 bits (18), Expect = 0.91
Identities = 18/18 (100%), Gaps = 0/18 (0%)
Strand=Plus/Plus

```
Query 7          CATCAGTTACTGTGTTGA 24
          |||||
sbjct 20573499    CATCAGTTACTGTGTTGA 20573516
```

Features flanking this part of subject sequence:
[14909 bp at 5' side: hypothetical protein LOC202134](#)
[47766 bp at 3' side: hypothetical protein isoform 1](#)

Score = 30.2 bits (15), Expect = 56
Identities = 15/15 (100%), Gaps = 0/15 (0%)
Strand=Plus/Plus

```
Query 10         CAGTTACTGTGTTGA 24
          |||||
sbjct 20367234    CAGTTACTGTGTTGA 20367248
```

Features in this part of subject sequence:
[delta-sarcoglycan isoform 1](#)
[delta-sarcoglycan isoform 2](#)

Score = 28.2 bits (14), Expect = 221
Identities = 14/14 (100%), Gaps = 0/14 (0%)
Strand=Plus/Minus

```
Query 7          CATCAGTTACTGTG 20
          |||||
sbjct 958772      CATCAGTTACTGTG 958759
```

Features flanking this part of subject sequence:
[62771 bp at 5' side: hypothetical protein LOC408263](#)